

An efficient copper catalyzed 3D mesoporous aluminosilicate for the synthesis of dibenzodiazonines in Ullmann cross-coupling reaction

Udayakumar Veerabagu ^{a,b,*}, Gowsika Jaikumar ^c, Pandurangan Arumugam ^b, Sabarathinam Shanmugam ^d and Lu Fushen ^{a*}

^a Department of Chemistry and Key Laboratory for Preparation and Application of Ordered Structural Materials of Guangdong Province, Shantou University, Guangdong 515063, China

^b Department of Chemistry, Anna University, Guindy, Chennai 600025, India

^c Department of Chemistry, Pachaiyappa's College, University of Madras, Chennai 600030, India

^d Bioprocess and Biomaterials Laboratory, Department of Microbial Biotechnology, Bharathiar University, Coimbatore 641046, India

* Corresponding authors. E-mail: udaya_taurus@yahoo.co.in; fslu@stu.edu.cn

Figure Captions

NMR Spectra

¹ H and ¹³ C NMR spectra for 13H-dibenzo[b,e][1,4]diazonine	2-3
¹ H and ¹³ C NMR spectra for 13H-dibenzo[b,e][1,4]diazonin-3-ol	4-5
¹ H and ¹³ C NMR spectra for 2,3-dimethyl-13H-dibenzo[b,e][1,4]diazonine	6-7
¹ H and ¹³ C NMR spectra for 3-bromo-13H-dibenzo[b,e][1,4]diazonine	8-9
¹ H and ¹³ C NMR spectra for 3-chloro-13H-dibenzo[b,e][1,4]diazonine	10-11



















