

Supplementary Information

Supramolecular Gel Assisted Synthesis of Co₂P Nanosheets as Efficient and Stable Catalyst for Oxygen Reduction Reaction

Zexing Wu^{a, 1,*}, Min Song^{a,1}, Jie Wang^{b,*} and Xien Liu^{a,*}

^a Key Laboratory of Sensor Analysis of Tumor Marker of Education Ministry; State Key Laboratory Base of Eco-chemical Engineering, College of Chemistry and Molecular Engineering, Qingdao University of Science & Technology, 53 Zhengzhou Road, 266042, Qingdao P. R. China. E-mail: splswzx@qust.edu.cn; liuxien@qust.edu.cn

^b Department of Applied Physics, The Hong Kong Polytechnic University, Hung Horn, Kowloon, Hong Kong. E-mail: jw65111@gmail.com

* Corresponding author.




Figure S1 (a) XRD pattern of Co_2P without carbon matrix. (b) LSV curve of Co_2P .




Figure S2 SEM image of $\text{Co}_2\text{P}/\text{C}$ (a) and corresponding Co (b), P (c) and C (d) elemental mappings.




Figure S3 EDS result of Co₂P/C.





Figure S4 TEM images of Co₂P/C after durability test.




Figure S5 PXRD of Co₂P/C after durability test.