Nano-assemblies of cationic mPEG brush block copolymers with gadolinium polytungstate [Gd(W₅O₁₈)₂]⁹⁻ form stable, high relaxivity MRI contrast agents

Joanne Ly,^{a,b} Yuhuan Li, ^{a,b} Mai N. Vu,^{a,b} Bradford A. Moffat^c, Kevin S. Jack^d, John F. Quinn,^{a,b} Michael R. Whittaker^{a,b} and Thomas P. Davis^{a,b,e}

 ^a ARC Centre of Excellence in Convergent Bio-Nano Science & Technology, Monash Institute of Pharmaceutical Sciences, Monash University, 381 Royal Parade, Parkville, Victoria 3052, Australia.
^b Drug Delivery, Disposition and Dynamics, Monash Institute of Pharmaceutical Science, Monash

University, 381 Royal Parade, Parkville, VIC 3052, Australia

^c Melbourne Brain Centre Imaging Unit, Department of Anatomy and Neuroscience, The University of Melbourne, Parkville 3010 Victoria, Australia,

^d Centre for Microscopy and Microanalysis, The University of Queensland, St Lucia, QLD 4072, Australia

^e Department of Chemistry, University of Warwick, Gibbet Hill, Coventry, ULCV4 7AL, United Kingdom



Figure S1. XPS measurements of GdWO with survey scan (a) and high resolution scan of Na, O, Gd and W ions (b). There is a higher amount of oxygen atoms than the theoretical amount due to the Auger effect of the sodium atom.



Figure S2. Thermogravimetric analysis of GdWO



Figure S3. UV-Vis spectra of GdWO from pH 5.9 – pH 9.0





Figure S4. GPC molecular weight distribution of PEG macroRAFT agent and PDMAEMA₂₀ and PDMAEMA₄₀ (a). After chain extension of the macroRAFT, PDMAEMA₂₀ and PDMAEMA₄₀ shift to the right (towards higher molecular weight) indicating successful addition of the DMAEMA block. The ¹H NMR spectra of PEG macroRAFT (b), PDMAEMA₂₀ (c) and PDMAEMA₄₀ (d).



Figure S5. DLS determined number average particle size of polymer PDMAEMA₂₀ (a) and PDMAEMA₄₀ (b) in aqueous solution at pH 7.4 and pH 6.0.



Figure S6. Thermogravimetric analysis of nano-assembly ^{0.2}P₂₀@GdWO before and after dialysis.



Figure S7. DLS determined the number average particle size of the P_{40} @GdWO nano-assemblies at week 1 and week 7 (a) and ζ -potential of the nano-assemblies (b).



Figure S8. TEM images of the ^{0.3}P₂₀@GdWO (a), ^{0.7}P₂₀@GdWO (b) and ^{0.9}P₂₀@GdWO (c).



Figure S9. Cytotoxicity study of the P20@GdWO nano-assemblies with AlamarBlue assay