

Electronic Supplementary Information (ESI) for

Magnolol dimer-derived fragments as PPAR γ -selective probes

Dominik Dreier,^a Mirta Resetar,^b Veronika Temml,^c Lukas Rycek,^a Nicolas Kratena,^a Michael Schnürch,^a Daniela Schuster,^d Verena M. Dirsch^b and Marko D. Mihovilovic*^a

^a Institute of Applied Synthetic Chemistry, TU Wien, Getreidemarkt 9/163, A-1060 Vienna, Austria. E-mail: marko.mihovilovic@tuwien.ac.at



^b Department of Pharmacognosy, University of Vienna, Althanstraße 14, 1090 Vienna, Austria



^c Institute of Pharmacy/Pharmacognosy and Center for Molecular Biosciences Innsbruck, University of Innsbruck, Innrain 80/82, 6020 Innsbruck, Austria



^d Institute of Pharmacy/Pharmaceutical Chemistry and Center for Molecular Biosciences Innsbruck, University of Innsbruck, Innrain 80/82, 6020 Innsbruck, Austria; Paracelsus Medical University Salzburg, Institute of Pharmacy, Department of Pharmaceutical and Medicinal Chemistry, Strubergasse 21, 5020 Salzburg, Austria.



Table of contents



NMR spectra of 2	2
NMR spectra of 3	3
NMR spectra of 4	4
NMR spectra of 6	5
NMR spectra of 7	6
NMR spectra of 8	7
NMR spectra of 9	8
NMR spectra of 13	9
NMR spectra of 14	10
NMR spectra of 15	11
NMR spectra of III	12
NMR spectra of IV	13
NMR spectra of V	14







































16

