

Electronic Supplementary Information

Bicyclo[2.2.1]heptane Containing N,N'-diarylsquaramides CXCR2 Selective Antagonists as Anti-cancer Metastasis Agents

Jin-Xin Che,^a Zhi-Long Wang,^b Xiao-Wu Dong,^a You-Hong Hu,^b Xin Xie^{*b, c} and
Yong-Zhou Hu^{*a}

^a ZJU-ENS Joint Laboratory of Medicinal Chemistry, College of Pharmaceutical
Sciences, Zhejiang University, Hangzhou, PR China.

^b State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica,
Chinese Academy of Sciences, Shanghai, PR China

^c CAS Key Laboratory of Receptor Research, the National Center for Drug Screening,
Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, PR
China.

*Corresponding authors:


E-mail: huyz@zju.edu.cn (Yong-Zhou Hu) ; E-mail: xxie@mail.shnc.ac.cn (Xin Xie)

Table of contents

Figure S1. CXCR2 homologous modeling	2
Table S1. CXCR1 and CXCR2 IC ₅₀ curve graphs.....	3
NMR spectras of compound 1a and 2e	5

Figure S1. CXCR2 homologous modeling

Align Sequence to Templates



Homologous Modeling (Build Homology Models, Standard Dynamic Cascade)









Table S1. CXCR1 and CXCR2 IC₅₀ curve graphs





1f





2a





2b





2c





2d





2e





2f



2g



2h



NMR spectras of compound 1a and 2e

