Yunnanterpene G, a Spiro-triterpene from the Roots of *Cimicifuga foetida* Down-regulates the Expressions of CD147 and MMPs in PMA Differentiated THP-1 Cells

Ni-Hong Lu,^a Zi-Wei Zhang,^b Rui-Wei Guo,^b Li-Xia Yang,^{b,*} Ya-Xian Song,^a Jin-Shan Ye,^b Yan-Kun Shi^b

^aDepartment of Postgraduate, Kunming Medical University, Yunnan, 650500, China

^bDepartment of Cardiology, Kunming General Hospital of Chengdu Military Area, Yunnan, 650032, China

Supporting Information List

Page 3-12, ¹H, ¹³C, HSQC, HMBC, COSY and ROESY NMR spectra, UV and IR spectra, and OR and HREIMS experiments of compound **1**.

Page 13, **Figure S11.** Cell morphology and expression of CD68 and CD11b of differentiated and undifferentiated THP-1 cells

Page 14, Table S1. Primers' information for CD147 and MMPs in the present study

Figure S1. ¹H (100 Hz) NMR Spectrum of Yunnanterpene G (1) in Pyridine- d_5



Figure S2. ¹³C (150 Hz) NMR Spectrum of Yunnanterpene G (1) in Pyridine- d_5





Figure S3. HSQC (600 Hz) Spectrum of Yunnanterpene G (1) in Pyridine- d_5





Figure S4. HMBC (600 Hz) Spectrum of Yunnanterpene G (1) in Pyridine- d_5

(Indd) []



Figure S5. ¹H-¹H COSY (600 Hz) Spectrum of Yunnanterpene G (1) in Pyridine-*d*₅

(II (ppu)



Figure S6. ROESY (600 Hz) Spectrum of Yunnanterpene G (1) in Pyridine- d_5

Figure S7. HREIMS of Yunnanterpene G (1)

Page 1

Autospec Premier P776 1

Elemental Composition Report

Single Mass Analysis Tolerance = 10.0 PPM / DBE: min = -10.0, max = 120.0 Selected filters: None

Monoisotopic Mass, Odd and Even Electron Ions 22 formula(e) evaluated with 1 results within limits (up to 51 closest results for each mass) Elements Used: C: 0-200 H: 0-400 O: 5-7 Na: 1-1 wgq-9 KIB 15:16:29 20-Apr-2016 M160420E-05 15 (1.103) Magnet El+

500.00 500.10 500.20 500.30 m/z 0-499.40 • • • 499.50 499.60 499.70 499.80 499.90 Minimum: Maximum: -10.0 120.0 200.0 10.0 Calc. Mass mDa PPM DBE Mass Formula 523.3029 523.3036 C30 H44 O6 Na -0.7 -1.3 8.5



Figure S8. UV spectrum of Yunnanterpene G (1)



Figure S9. IR spectrum of Yunnanterpene G (1)

			1.1.1			1.		
Optica	I rotation m	easurement						
Model No.	: P-1020 (A Sample	060460638) Mode	Data					
	, and the second	Mode	Dala	Blank	Temp. Cell Temp Point	Date Comment Sample Name	Light Filter Operator	Cycle Time Integ Time
No.1	21 (1/3)	Sp.Rot	-40.0000	-0.0044 0.0000	19.2 10.00	Tue Oct 13 19:48:43 2015 0.00110g/mL MeOH	Na 589nm	2 sec 10 sec
No.2	21 (2/3)	Sp.Rot	-40.9090	-0.0045 0.0000	19.1 10.00	KYS35 Tue Oct 13 19:48:56 2015 0.00110g/mL MeOH	Na 589nm	2 sec 10 sec - 39. 0909
No.3	21 (3/3)	Sp.Rot	-36.3640	-0.0040 0.0000	19.1 10.00 Cell	Tue Oct 13 19:49:10 2015 0.00110g/mL MeOH KYS35	Na 589nm	2 sec 10 sec
						ei in the second se		
				10	14			

Figure S10. $[a]_D$ value of Yunnanterpene G (1)



Figure S11. Cell morphology and expression of CD68 and CD11b of differentiated and undifferentiated THP-1 cells. **A.** Undifferentiated THP-1 cells; **B.** Induction of monocyte-macrophage differentiation by 100 nM PMA for 24 h; **C.** CD68 expression of the differentiated cells; **D.** CD11b expression of the differentiated cells. Expression of CD68 and CD11b were measured by flow cytometry using FITC-labeled anti-CD68 mAb and anti-CD11b mAb. Analysis was conducted on a PARTEC brand flow cytometer.

CTGGTACAAGATCACTGAC	EMMPRIN(H)_E
CIOCIACAROAICACIÓAC	
GAGGAACTCACGAAGAAC	EMMPRIN(H)-R
CTGAAGGACACACTAAAGA	MMP2(H)-F
CGATGGTATTCTGGTCAA	MMP2(H)-R
GGCAGATTCCAAACCTTT	MMP9(H)-F
GCAAGTCTTCCGAGTAGT	MMP9(H)-R
TCCATCAACACTGCCTAC	MMP14(H)-F
CCTCATCAAACACCCAATG	MMP14(H)-R
AAAGGGTCATCATCTCTG	GAPDH-F
GCTGTTGTCATACTTCTC	GAPDH-R

Table S1. Primers'	information f	for CD147	and MMPs in	the present study