

SUPPLEMENTARY DATA

Isolation of maltol derivatives from *Stellera chamaejasme* and the anti-atopic properties of maltol on skin lesions in DNCB-stimulated mice

Beom-Geun Jo,^{‡a} No-June Park,^{‡b} Su-Nam Kim,^b Jonghwan Jegal,^a Sangho Choi,^c Sang Woo Lee,^c Li Wan Yi,^d Seoung Rak Lee,^e Ki Hyun Kim^{*e} and Min Hye Yang^{*a}

^aCollege of Pharmacy, Pusan National University, Busan 46241, South Korea. E-mail: mhyang@pusan.ac.kr; Tel: +82-51-510-2811; +82-51-513-6754

^bNatural Products Research Institute, Korea Institute of Science and Technology, Gangneung 25451, South Korea

^cInternational Biological Material Research Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon 34141, South Korea


^dInstitute of Medicinal Plants, Yunnan Academy of Agricultural Sciences, Yunnan 650205, China

^eNatural Product Research Laboratory, School of Pharmacy, Sungkyunkwan University, Suwon, 16419, South Korea. E-mail: khkim83@skku.edu; Tel: +82-31-290-7700; Fax: +82-31-290-7730


Supporting Information Contents:

- S1.** The ¹H NMR spectrum of compound 3
- S2.** The ¹³C NMR spectrum of compound 3
- S3.** The ¹H-¹H COSY spectrum of compound 3
- S4.** The ¹H-¹H NOESY spectrum of compound 3
- S5.** The HMQC spectrum of compound 3
- S6.** The HMBC spectrum of compound 3


S1. The ^1H NMR spectrum of compound **3**




S2. The ^{13}C NMR spectrum of compound 3




S3. The ^1H - ^1H COSY spectrum of compound 3



S4. The ^1H - ^1H NOESY spectrum of compound **3**



S5. The HMQC spectrum of compound 3



S6. The HMBC spectrum of compound 3

