

Supporting Information

Electrocatalytic Water Oxidation over AlFe_2B_2

Dallas K. Mann,^{a,#} Junyuan Xu,^{b,#} Natalia E. Mordvinova,^c Vincent Yannello,^a Yasmine Ziouani,^b Noelia González-Ballesteros,^d Juliana P. S. Sousa,^b Oleg I. Lebedev,^c Yury V. Kolen'ko,^{b,*} and Michael Shatruk^{a,*}

^a Department of Chemistry and Biochemistry, Florida State University, 95 Chieftan Way, Tallahassee, FL 32306, USA

^b International Iberian Nanotechnology Laboratory, Braga 4715-330, Portugal

^c Laboratoire CRISMAT, UMR 6508, CNRS-Ensicaen, Caen 14050, France

^d Inorganic Chemistry Department, Biomedical Research Centre (CINBIO), Universidade de Vigo, Vigo 36210, Spain

E-mail: shatruk@chem.fsu.edu; yury.kolenko@inl.int

Contents	Page
Figure S1. Current density vs. applied potential curves normalized by the surface area	S2
Figure S2. EDX spectra of the AlFe_2B_2 sample subjected to 20 cycles of electrocatalytic activation.....	S2




Figure S1. Current density vs. applied potential curves measured for different OER electrocatalysts in a 1M KOH electrolyte solution and normalized by the BET surface area of the catalyst.




Figure S2. EDX spectra of the AlFe_2B_2 sample subjected to 20 cycles of electrocatalytic activation in a 1-M KOH electrolyte solution. The absence of any sizeable amount of Ni in the sample is obvious.