

Electronic Supporting Information

**Aromatic Identity, Electronic Substitution, and Sequence in
Amphiphilic Tripeptide Self-Assembly**

Jugal Kishore Sahoo¹, Calvin Nazareth¹, Michael A. VandenBerg¹, Matthew J. Webber<sup>*1, 2, 3,
4, 5, 6</sup>

1. Department of Chemical and Biomolecular Engineering, University of Notre Dame, Notre Dame, IN 46556, USA
2. Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA
3. Harper Cancer Research Institute, University of Notre Dame, Notre Dame, IN 46556, USA
4. Advanced Diagnostics and Therapeutics, University of Notre Dame, Notre Dame, IN 46556, USA
5. Warren Family Center for Drug Discovery and Development, University of Notre Dame, Notre Dame, IN 46556, USA
6. Center for Nanoscience and Technology (NDnano), University of Notre Dame, Notre Dame, IN 46556, USA

Correspondence should be addressed to
Prof. Matthew. J. Webber
Department of Chemical and Biomolecular Engineering
University of Notre Dame
205B McCourtney Hall
Notre Dame, IN, 46556 USA
E-Mail: mwebber@nd.edu

Contents

1. Chemical Characterisation of Tripeptides (^1H NMR, Mass Spec, and HPLC)
2. Additional Supporting Data
3. References

1. Chemical Characterisation of Tripeptides




Figure S1: ¹H NMR spectrum of Ac-FID-NH₂




Figure S2: Analytical HPLC chromatogram of Ac-FID-NH₂




Figure S3: Mass spectroscopy of Ac-FID-NH₂




Figure S4: ^1H NMR spectrum of Ac-YID-NH₂




Figure S5: Analytical HPLC chromatogram of Ac-YID-NH₂




Figure S6: Mass spectroscopy of Ac-YID-NH₂




Figure S7: ^1H NMR spectrum of Ac-WID-NH₂




Figure S8: Analytical HPLC chromatogram of Ac-WID-NH₂




Figure S9: Mass spectroscopy of Ac-WID-NH₂




Figure S10: ^1H NMR spectrum of Ac-IFD-NH₂




Figure S11: Analytical HPLC chromatogram of Ac-IFD-NH₂




Figure S12: Mass spectroscopy of Ac-IFD-NH₂




Figure S13: ^1H NMR spectrum of Ac-DIF-NH₂




Figure S14: Analytical HPLC of Ac-DIF-NH₂




Figure S15: Mass spectroscopy of Ac-DIF-NH₂




Figure S16: ^1H NMR spectroscopy of Ac-DFI-NH₂




Figure S17: Analytical HPLC of Ac-DFI-NH₂




Figure S18: Mass spectroscopy of Ac-DFI-NH₂




Figure S19: ¹H NMR spectrum of Ac-IDF-NH₂




Figure S20: Analytical HPLC of Ac-IDF-NH₂




Figure S21: Mass spectroscopy of Ac-IDF-NH₂




Figure S22: ^1H NMR spectrum of Ac-FDI-NH₂




Figure S23: Analytical HPLC of Ac-FDI-NH₂




Figure S24: Mass spectroscopy of Ac-FDI-NH₂




Figure S25: ^1H NMR spectrum of Ac-F_FID-NH₂




Figure S26: Analytical HPLC of Ac-F_EID-NH₂




Figure S27: Mass spectroscopy of Ac-F_FID-NH₂




Figure S28: ¹H NMR spectrum of Ac-F_IID-NH₂




Figure S29: Analytical HPLC of Ac-F₁ID-NH₂




Figure S30: Mass spectroscopy of Ac-F₁ID-NH₂




Figure S31: ^1H NMR spectrum of Ac-F_{NO₂}ID-NH₂




Figure S32: Analytical HPLC of Ac-F_{NO₂}ID-NH₂




Figure S33: Mass spectroscopy of $\text{Ac-F}_{\text{NO}_2}\text{ID-NH}_2$




Figure S34: ^1H NMR spectrum of $\text{Ac-F}_{\text{PO}_4}\text{ID-NH}_2$




Figure S35: Analytical HPLC of Ac-F_{PO4}ID-NH₂




Figure S36: Mass spectroscopy of Ac-F_{PO4}ID-NH₂

2. Additional Supporting Data




Figure S37: A, B) Oscillating rheology frequency sweep of FID and YID respectively.




Figure S38: Additional TEM images of the tripeptides: A and D) FID; B and E) YID; C and F) WID.




Figure S39: A, B, C) Oscillating rheology frequency sweep of F_FID , F_ID and $\text{F}_{\text{NO}_2}\text{ID}$ respectively.





Figure S40: Additional TEM images of the tripeptides at different magnifications: A and E) $\text{F}_{\text{PO}_4}\text{ID}$; B and F) F_FID ; C and G) F_ID ; D and H) $\text{F}_{\text{NO}_2}\text{ID}$.



Substitution (X)	Hammett Values
-OH	-0.37
-OPO ₃ H ₂	0.00
-H	0.00
-F	0.06
-I	0.18
-NO ₂	0.78

Table S1: Substitution groups and their corresponding Hammett σ_P Values (From Ref S1)




Figure S41: A, B, C) oscillating rheology frequency sweep of IFD, DIF and DFI respectively.




Figure S42: Additional TEM images of the tripeptides at different magnifications: A and F) IFD; B and G) DIF; C and H) FDI; D and I) IDF; E and J) FDI.

4. References:

S1. C. Hansch, A. Leo and R. W. Taft, *Chem. Rev.*, 1991, **91**, 165–195.