Carboplatin prodrug conjugated Fe_3O_4 nanoparticles for magnetically targeted drug delivery in ovarian cancer cells

Haiqin Song ^{a, 1}, Feifei Quan ^{b, 1}, Zhiqiang Yu ^c, Minhua Zheng ^a, Yan Ma ^b*, Haihua Xiao ^d*, Fang Ding ^e*

^a Department of General Surgery, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, and Shanghai Minimally Invasive Surgery Center, Shanghai, 200025 P. R. China.

^b The First Affiliated Hospital of University of South China, Hengyang, 421001, China. *Email: <u>nhfsmy@126.com</u>
^c School of Pharmaceutical Sciences, Guangdong Provincial Key Laboratory of New Drug Screening, Southern Medical University, Guangzhou, 510515, China

^d Beijing National Laboratory for Molecular Sciences, State Key Laboratory of Polymer Physics and Chemistry, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, P. R. China. *Email: hhxiao@iccas.ac.cn

^e Nanshan District Key Lab for Biopolymers and Safety Evaluation, Shenzhen Key Laboratory of Polymer Science and Technology, Guangdong Research Center for Interfacial Engineering of Functional Materials, College of Materials Science and Engineering, Shenzhen University, Shenzhen 518060, P. R. China. *Email:_ <u>dingfang0707@163.com</u>

¹ The two authors contributed equally to this work.



Figure S1. H&E staining analysis of major organs in the test mice injected with NPs@Carboplatin. Mice without any treatment were set as control. Mice were sacrificed at day 7, 14 and 30 post injection.