

Supporting Information

**Efficient enantiorecognition of amino acids under a stimuli-responsive system: Synthesis, characterization and application of electroactive rotaxane**

Datong Wu,<sup>a</sup> Fei Pan,<sup>a</sup> Gao-Chao Fan,<sup>b</sup> Ziming Zhu,<sup>c</sup> Li Gao,<sup>a</sup> Yongxin Tao<sup>a</sup> and


Yong Kong<sup>\*a</sup>

*<sup>a</sup>Jiangsu Key Laboratory of Advanced Materials and Technology, School of Petrochemical Engineering, Changzhou University, Changzhou 213164, China.*

*\*Email: yzkongyong@cczu.edu.cn*

*<sup>b</sup>Key Laboratory of Optic-electric Sensing and Analytical Chemistry for Life Science, MOE, Shandong Key Laboratory of Biochemical Analysis, Qingdao University of Science and Technology, Qingdao 266042, China*

*<sup>c</sup>College of Life Science and Technology, China Pharmaceutical University, Nanjing 211198, China*



**GPC Results**

	Dist Name	Mn	Mw	MP	Mz	Mz+1	Mv	Polydispersity	% Area
1		5780	6481	5922	7239	7982		1.121290	100.00

**Fig. S1.** GPC results of polymer-2.