

SUPPLEMENTARY

Table 1. PAES concentration levels (ng/g) determined by liquid extraction (LE) and direct immersion microextraction (SPME) in specimens of *Danafungia scruposa* (DF) and *Tridacna maxima* (TM) collected along the Faafu atoll, Maldives. BDL = beyond the detection limit. Data are corrected by subtracting the average of five procedural blank

| Sample | Extraction method | MEHP | DMP | DEP | DBP | BBzP | DEHP |
|---------------|--------------------------|-------------|------------|------------|------------|-------------|-------------|
| DF1 | LE | BDL | BDL | BDL | 1,8 | BDL | 2,3 |
| | SPME | BDL | BDL | 0,4 | 1,7 | BDL | 2,2 |
| DF2 | LE | BDL | 1,6 | BDL | 1,2 | BDL | 4,6 |
| | SPME | BDL | 1,4 | BDL | 0,9 | BDL | 4,5 |
| DF3 | LE | BDL | BDL | BDL | 1,4 | 1 | 2,5 |
| | SPME | BDL | BDL | BDL | 1,2 | 0,9 | 2,5 |
| TM1 | LE | 1,8 | 4,5 | 2,3 | 2,4 | 1,9 | 7,4 |
| | SPME | 1,7 | 4,3 | 2,1 | 2,0 | 1,7 | 6,7 |