

Electronic Supplementary Information (ESI) for Biomaterials Science

Effective systemic siRNA delivery using dual-layer protected long-circulating nanohydrogel containing inorganic core

Yue Yin^{a,1}, Min Sang Lee^{a,1}, Jung Eun Lee^a, Su Yeon Lim^a, E Seul Kim^a, Jaeback Jeong^a,
Dahwun Kim^a, Jaeyun Kim^b, Doo Sung Lee^{b,*}, and Ji Hoon Jeong^{a,*}

^aSchool of Pharmacy, Theranostic Macromolecules Research Center, Sungkyunkwan University,
Suwon 440-746, Republic of Korea

^bSchool of Chemical Engineering, Theranostic Macromolecules Research Center,
Sungkyunkwan University, Suwon 440-746, Republic of Korea

*Corresponding authors:

Doo Sung Lee, Ph.D.

Tel.: +82-31-299-6851; Fax: +82-31-299-6857; e-mail: dslee@skku.edu

Ji Hoon Jeong, Ph.D.

Tel.: +82-31-290-7783; Fax: +82-31-292-8800; e-mail: jhjeong@skku.edu

¹The first two authors contributed equally to this study.

Supplementary result




Figure S1. ¹H-NMR spectrum of PEG-dextran-dopa.