

SUPPLEMENTARY INFORMATION

Water-Mediated Efficient Synthesis of Bismuth Oxyiodide for Several Distinct Morphologies

Yang Long^a, Siyuan Tan^a, Qiang Han^a, Yongjian Ai^a, Yuqiang Sheng^a, Yi Wang^{*b}, Haibo Wang^c,
Youliang Xie^d, Qionglin Liang^{*a}, Mingyu Ding^{*a}

a: MOE Key Laboratory of Bioorganic Phosphorus Chemistry & Chemical Biology; Department of Chemistry, Center for Synthetic and Systems Biology, Tsinghua University, Beijing, 100089, P. R. China.

b: Key Laboratory of Marine Environmental Corrosion and Bio-fouling, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, 266071, P. R. China.

c: School of Pharmaceutical sciences, Liaoning University of Traditional Chinese Medicine, Dalian 116600, P. R. China.

d: Mathematical Engineering Academy of Chinese Medicine, Guangzhou University of Chinese Medicine, Guangzhou 510006, P. R. China.




Figure S1, XRD patterns of the products prepared without reaction water.




Figure S2, SEM image of the products prepared at 2.5 mL of reaction water.




Figure S3, EDS analysis of bowl-like BiOI microstructures