

Resonant photoionization of O₂ up to the fourth ionization threshold: Supporting Information

Sonia Marggi Poullain,^a Markus Klinker,^a Jesús González-Vázquez,^a and Fernando Martín^{a,b,c}

1 Partial Photoionization Cross Section

The partial photoionization cross section into the a⁴Π_u, A²Π_u and b⁴Σ_g⁻ ionic states of O₂⁺ are depicted in Figures 1, 2 and 3, respectively.




Figure 1: Computed Partial Cross section into O₂⁺ (a⁴Π_u).

^{0a} Departamento de Química, Módulo 13, Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Madrid, Spain.

^{0b} Instituto Madrileño de Estudios Avanzados en Nanociencia (IMDEA-Nanociencia), Cantoblanco, 28049 Madrid, Spain.

^{0c} Condensed Matter Physics Center (IFIMAC), Universidad Autónoma de Madrid, 28049 Madrid, Spain.




Figure 2: Computed Partial Cross section into O_2^+ ($A^2\Pi_u$).




Figure 3: Computed Partial Cross section into O_2^+ ($b^4\Sigma_g^-$).

2 Total Photoionization Cross Sections

The computed total photoionization cross section for the ${}^3\Sigma_u^-$ and ${}^3\Pi_u$ symmetries are shown in Fig. 4.




Figure 4: Total photoionization cross section computed for the ${}^3\Sigma_u^-$ (orange, labelled Z axis) and ${}^3\Pi_u$ (violet, labelled X axis) total symmetries.