

Light-stable Polypyridine Silver(I) Complexes of 1,3,5-Triaza-7-phosphaadamantane (PTA) and 1,3,5-Triaza-7-phosphaadamantane-7-sulfide (PTA=S): Significant Antiproliferative Activity of Representative Examples in Aqua Media

Sabina W. Jaros^a, Urszula Śliwińska-Hill^b, Agata Białońska^a, Dmytro S. Nesterov^c, Jerzy Sokolnicki^a, Piotr Kuroпка^d, Barbara Bażanów^b and Piotr Smoleński^{*,a}

^a Faculty of Chemistry, University of Wrocław, F. Joliot-Curie 14, 50-383 Wrocław, Poland

^b Department of Analytical Chemistry, Faculty of Pharmacy, Wrocław Medical University, Borowska 211 A, 50-566 Wrocław, Poland

^c Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

^b Department of Veterinary Microbiology, Wrocław University of Environmental and Life Sciences, Norwida 31, 50-375 Wrocław, Poland

^d Department of Histology and Embryology, Wrocław University of Environmental and Life Sciences, Norwida 31, 50-375 Wrocław, Poland

*Corresponding author.

E-mail addresses: piotr.smolenski@chem.uni.wroc.pl (P. Smoleński).




Figure S1. Image of the A549 cells after 4 hours incubation in volume 1000 μ l of DMEM. Visible single in group 1 (arrow) and multiple death or dying cells in group 3. DAPI and propidium iodide Mag 200x.




Figure S2. Morphometrical analysis of live/death cell ratio in A549 cell line after 4 hours of incubation in presence of **1** and **3**. The differences were statistically significant at $p \leq 0,05$.




Figure S3. Morphometrical analysis of live/death cell ratio in A549 cell line after 12 hours of incubation at different concentrations of **1** and **3**. The differences were statistically significant at $p \leq 0,05$.