

Supporting information

Stainless steel mesh-GO/Pd NPs: catalytic application for Suzuki-Miyaura and Stille coupling reactions in eco-friendly media

**Bahareh Feizi Mohazzab,^a Babak Jaleh^{a,*}, Zahra Issaabadi,^b Mahmoud Nasrollahzadeh^{b,*}
and Rajender S. Varma^{c,*}**

^aDepartment of Physics, Faculty of Science, Bu-Ali Sina University, Postal Code 65174 Hamedan, Iran. E-mail:

jaleh@basu.ac.ir


^bDepartment of Chemistry, Faculty of Science, University of Qom, Qom 3716146611, Iran. E-mail:

mahmoudnasr81@gmail.com; Fax: +98 25 32103595; Tel: +98 25 32850953

^cRegional Centre of Advanced Technologies and Materials, Department of Physical Chemistry, Faculty of Science,

Palacky University, Šlechtitelů 27, 783 71 Olomouc, Czech Republic. E-mails: rajvarma@hotmail.com;

Varma.Rajender@epa.gov; Fax: +1 (513)-569-7677; Tel.: +1 (513)-487-2701







Figure S1. ¹H NMR spectrum (500 MHz, CDCl₃) of biphenyl.







Figure S2. ¹H NMR spectrum (500 MHz, CDCl₃) of 4-methoxybiphenyl.







Figure S3. ¹H NMR spectrum (500 MHz, CDCl₃) of 4-nitrobiphenyl.







Figure S4. ^1H NMR spectrum (500 MHz, CDCl_3) of 1,1'-biphenyl-4-carbonitrile.






Figure S5. ¹H NMR spectrum (500 MHz, CDCl₃) of 4-aminobiphenyl