

Synthesis and evaluation of novel 2,4-disubstituted arylthiazole derivatives against *T.brucei*


Markos-Orestis Georgiadis,^a Violeta Kourbeli,^a Ioannis P. Papanastasiou,*^a Andrew Tsotinis,^a Martin C. Taylor ^b and John M. Kelly ^b



^aSchool of Health Sciences, Department of Pharmacy, Division of Pharmaceutical Chemistry, National and Kapodistrian University of Athens, Panepistimioupoli-Zografou, 157 84 Athens, Greece


^bDepartment of Pathogen Molecular Biology, London School of Hygiene and Tropical Medicine, Keppel Street, London

Supporting Information


¹HNMR and ¹³C NMR spectra









¹³C NMR (150 MHz, DMSO-*d*₆) of 2-{2-[3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl]thiazol-4-yl}N-methylethan-1-amine dihydrochloride (**1b**)










^1H NMR (400 MHz, CDCl_3) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl]thiazol-4-yl}ethyl}acetamide (**1d**)




^{13}C NMR (50 MHz, CDCl_3) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl]thiazol-4-yl}ethyl}acetamide (**1d**)








¹H NMR (400 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}-4-fluorobenzamide (**1g**)




¹³C NMR (50 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}-4-fluorobenzamide (**1g**)

MG-70B






¹H NMR (400 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}-3-methylfuran-2-ylcarboxamide (**1h**)

MG-70C
13C dec
ns 1000




¹³C NMR (50 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}-3-methylfuran-2-ylcarboxamide (**1h**)










¹H NMR (400 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}piperidine-1-ylcarboxamide (**1j**)




¹³C NMR (150 MHz, CDCl₃) of *N*-{2-[2-(3-(1-tricyclo[3.3.1.1^{3,7}]decyl)-4-fluorophenyl)thiazol-4-yl]ethyl}piperidine-1-ylcarboxamide (**1j**)









¹H NMR (400 MHz, DMSO-*d*₆) of 2-{2-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-4-yl}ethan-1-amine dihydrochloride (**2a**)





¹³C NMR (75 MHz, DMSO-*d*₆) of 2-{2-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-4-yl}ethan-1-amine dihydrochloride (**2a**)







¹H NMR (600 MHz, DMSO-*d*₆) of 2-{2-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-4-yl}-*N*-methylethan-1-amine difumarate (**2b**)







¹³C NMR (150 MHz, DMSO-*d*₆) of 2-{2-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-4-yl}-*N*-methylethan-1-amine difumarate (**2b**)








































^a¹H NMR (400 MHz, DMSO-*d*₆) of 2-{4-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-2-yl}-*N,N*-dimethylethan-1-amine dihydrochloride (**3d**)




^a¹³C NMR (150 MHz, DMSO-*d*₆) of 2-{4-[4-(1-tricyclo[3.3.1.1^{3,7}]decyl)phenyl]thiazol-2-yl}-*N,N*-dimethylethan-1-amine dihydrochloride (**3d**)







¹H NMR (400 MHz, DMSO-*d*₆) of 1-((1*R*,4*R*)-7,7-Dimethyl-2-oxobicyclo[2.2.1]hept-1-yl)-*N*-(2-(2-phenylthiazol-4-yl)ethyl)methanesulfonamide hydrochloride (**4b**)



¹³C NMR (150 MHz, DMSO-*d*₆) of 1-((1*R*,4*R*)-7,7-Dimethyl-2-oxobicyclo[2.2.1]hept-1-yl)-*N*-(2-(2-phenylthiazol-4-yl)ethyl)methanesulfonamide hydrochloride (**4b**)

