

ELECTRONIC SUPPLEMENTARY MATERIALS

Copper(II) Complexes with Alloferon Analogues Containing Phenylalanine H6F and H12F Stability and Biological Activity

Lower Stabilization of Complexes Compared to Analogues Containing Tryptophan

**Teresa Kowalik-Jankowska^{a*}, Monika Lesiów^a, Katarzyna Krupa^a, Mariola Kuczer^a,
Elżbieta Czarniewska^b**

*^a Faculty of Chemistry, University of Wrocław, 14 F. Joliot-Curie Str., 50-383 Wrocław,
Poland*

*^b Department of Animal Physiology and Development, Institute of Experimental Biology,
Adam Mickiewicz University, 89 Umultowska Str., 61-614 Poznań, Poland*

*corresponding author: teresa.kowalik-jankowska@chem.uni.wroc.pl




Figure S1. Species distribution diagram for Cu(II) complexes with Allo12F. Charges are omitted for clarity. $[\text{L}] = 0.001 \text{ M}$, M/L molar ratio = 1:1




Figure S2. Species distribution diagram for Cu(II) complexes with Allo12F. Charges are omitted for clarity. $[\text{L}] = 0.001 \text{ M}$, M/L molar ratio = 2:1




Figure S3. Species distribution diagram for Cu(II) complexes with Allo12F. Charges are omitted for clarity. $[\text{L}] = 0.001 \text{ M}$, M/L molar ratio = 3:1