

Electronic Supplementary Information

A 9,9'-bifluorenylidene derivative containing four 1,1-dicyanomethylene-3-indanone end-capped groups as an electron acceptor for organic photovoltaic cells

Tianzhi Yu^{1,*}, Qianguang Yang¹, Tong Zhang¹, Yuling Zhao², Chengjin Wei¹, Yanmei Li¹,

Hui Zhang¹, Dingqin Hu^{3,*}

¹Key Laboratory of Opto-Electronic Technology and Intelligent Control (Ministry of Education), Lanzhou Jiaotong University, Lanzhou 730070, China

²School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, China

³Organic Semiconductor Research Center, Chongqing Institute of Green and Intelligent Technology, Chinese Academy of Sciences, Chongqing 400714, China

*Corresponding Author. E-mail: yutzh@mail.lzjtu.cn (T.Z. Yu); hudingqin@cigit.ac.cn (D. Q. Hu)




Fig. S1. ¹H NMR spectrum of the compound I




Fig. S2. ¹³C NMR spectrum of the compound I





Fig. S3. Mass spectrum of the compound I

Fig. S4. ¹H NMR spectrum of the compound II




Fig. S5. ^{13}C NMR spectrum of the compound II




Fig. S6. Mass spectrum of the compound II




Fig. S7. ¹H NMR spectrum of the compound **III**




Fig. S8. ¹³C NMR spectrum of the compound **III**





Fig. S9. Mass spectrum of the compound III

Fig. S10. ¹H NMR spectrum of the compound IV




Fig. S11. ^{13}C NMR spectrum of the compound IV




Fig. S12. Mass spectrum of the compound IV





Fig. S14. ¹³C NMR spectrum of the compound **BF-TDCI₄**




Fig. S15. Mass spectrum of the compound **BF-TDCI₄**