

Anti-pseudo-allergic capacity of alkaloids screened from *Uncaria Rhynchophylla*

Yitong Xie‡, Di Wei‡, Tian Hu, Yajing Hou, Yuanyuan Lin, Huaizhen He†, Cheng Wang†
School of Pharmacy of Pharmacy, Xi'an Jiaotong University, Xi'an 710061 China.

‡These two authors contribute equally.

†Correspondence Author: Cheng Wang and Huaizhen He

Address for correspondence

School of Pharmacy, Xi'an Jiaotong University, Yanta Westroad, Xi'an 710061, China

E-mail: hehuaizhen@mail.xjtu.edu.cn; chengwang@mail.xjtu.edu.cn.

Tel.: 86-29-82656788

Fax: 86-29-82655451




Fig. S1. Repeatability of ciprofloxacin on MRGPRX2/CMC column.






Fig. S2. Chromatogram of G1-G7 on MRGPRX2/CMC column.



G6



G7




Fig. S3. Mass spectra of G2-G7

Table.S1 HPLC-MS/MS information of compounds in UR

Peak No.	t _R (min)	Formula	Selected ion	Expected	Detected
2	11.12	C ₂₂ H ₂₆ N ₂ O ₄	[M+H] ⁺	383.45	383.30
3	12.88	C ₂₂ H ₂₈ N ₂ O ₄	[M+H] ⁺	385.46	385.30
4	14.44	C ₂₂ H ₂₆ N ₂ O ₄	[M+H] ⁺	383.45	383.30
5	18.86	C ₂₂ H ₂₈ N ₂ O ₄	[M+H] ⁺	385.46	385.35
6	20.76	C ₂₂ H ₂₆ N ₂ O ₃	[M+H] ⁺	367.46	367.25
7	26.73	C ₂₂ H ₂₈ N ₂ O ₃	[M+H] ⁺	369.46	369.30





Fig. S4.A The retained time and structures of G2-G7. B. Chromatogram of Rhy, Isorhy, Cor, Isocor, Hir, and Hie on MRGPRX2/CMC column.




Fig. S5. Effect of Rhy(a), Isorhy(b), Cor(c), Isocor(d), Hirsutine(e), and Hirsuteine(f) on cell viability on LAD2 cells measured by CCK-8 assay. Data are presented as the means \pm S.E.M (n=5); *P<0.05, **P<0.01 and ***P<0.001 vs. control cells.

Fig. 7. Rhy(a), Isorhy(b), Cor(c) and Isocor(d) degranulation effect on LAD2 mast cells. A. The β -Hexosaminidase release rate was analyzed after treatment of four alkaloid. B. The release level of histamine was determined after treatment of four alkaloid. Data are presented as the means \pm S.E.M (n=3); *P<0.05, **P<0.01 and ***P<0.001 vs. C48/80 group.




Figure S6. The Ca^{2+} influx of isorhy (a) and cor (b) in MRGPRX2 cells.