

## Supplementary Information

### **Sr-doped NiO<sub>3</sub> nanorods synthesized by simple sonochemical method as excellent materials for voltammetric determination of quercetin†**

Venkatachalam Vinothkumar,<sup>a</sup> Arumugam Sangili,<sup>a</sup> Shen-Ming Chen,<sup>\*a</sup> Pitchaimani Veerakumar<sup>\*ab</sup> and King-Chuen Lin<sup>\*ab</sup>

<sup>a</sup>Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, No. 1, Chung-Hsiao East Road, Section 3, Taipei 10608, Taiwan, ROC

<sup>b</sup>Department of Chemistry, National Taiwan University, No. 1, Roosevelt Road, Section 4, Taipei, 10617, Taiwan, ROC


<sup>c</sup>Institute of Atomic and Molecular Sciences, Academia Sinica, No. 1, Roosevelt Road, Section 4, Taipei, 10617, Taiwan, ROC

#### **\*Corresponding Authors**

E-mail address: smchen78@ms15.hinet.net (S.-M. Chen)

E-mail address: spveerakumar@gmail.com (P. Veerakumar)

E-mail address: kclin@ntu.edu.tw (K.-C. Lin)



**Fig. S1.** (a) Nyquist plot, (b) CV curves of 4.0, 6.0, and 8.0  $\mu\text{L}$  SNO NRs-modified GCE recorded in 5.0 mM  $[\text{Fe}(\text{CN})_6]^{3-/4-}/0.1$  M KCl.