

Supplemental Information

Enhancing Chemoradiation of Colorectal Cancer Through Targeted
Delivery of Raltitrexed by Hyaluronic Acid Coated Nanoparticles

Justin G. Rosch^a, Madeleine R. Landry^a, Charles R. Thomas, Jr.^b, and Conroy Sun^{a,b,*}

^a Department of Pharmaceutical Sciences, College of Pharmacy, Oregon State University, Portland, OR
97201, USA

^b Department of Radiation Medicine, School of Medicine, Oregon Health & Science University, Portland,
OR 97239, USA

*Corresponding Author:

Conroy Sun, Ph.D.

Department of Pharmaceutical Sciences

Oregon State University

2730 SW Moody Ave

Portland, OR 97201-5041, USA

Email: sunc@ohsu.edu

Phone: 503-346-4699




Figure S1. TEM of bare CML nanoparticles prior to layer-by-layer deposition.




Figure S2. Fluorescence microscopy images of no treatment and CML-PLA-HA treated wells after 24 h exposure. The no treatment well is the same as presented in **Figure 3**. The hyaluronic acid only nanoparticles showed similar fluorescence to the HARPs.




Figure S3. 24, 48, and 72 h viability curves for raltitrexed and HARPs.




Figure S4. Stacked histograms showing the changing fluorescent signal with different treatments (γ H2AX assay).