

Carbon Nitride Mediated Strong Metal-Support Interaction in Au/TiO₂ Catalyst for Aerobic Oxidative Desulfurization

Mingyang Liu ^a, Jing He ^a, Peiwen Wu ^a, Linjie Lu ^a, Chao Wang ^b, Linlin Chen ^a,
Mingqing Hua ^a, Wenshuai Zhu, *,^a and Huaming Li ^a

^a School of Chemistry and Chemical Engineering, Institute for Energy Research
Jiangsu University, Zhenjiang, Jiangsu, 212013, P. R. China

^b School of Environment and Safety Engineering, Jiangsu University, Zhenjiang,
Jiangsu, 212013, P. R. China

* E-mail: zhuws@ujs.edu.cn




Figure S1. TEM images of catalysts (a) Au-TiO₂@C₃N₄ (b) Au-TiO₂




Figure S2. Contact angle for an oil droplet on the surface of Au-TiO₂




Figure S3. Contact angle for an oil droplet on the surface of Au-TiO₂@C₃N₄




Figure S4. Contact angle for an oil droplet on the surface of Au-TiO₂-800




Figure S5. HRTEM image of Au-TiO₂@C₃N₄ catalyst




Figure S6. XRD patterns of (a) Au-TiO₂-800 (b) Au-TiO₂@C₃N₄-800




Figure S7. HRTEM images of Au-TiO₂@C₃N₄-800.




Figure S8. GC-MS analysis of oil phase and catalyst phase during the reaction