

Supporting Information

New *Leishmania donovani* nucleoside hydrolase inhibitors from Brazilian flora.

Charlotte Nirma,^a Gregorio Torres Rangel,^a Marina Amaral Alves,^{a,b} Livia Marques Casanova,^a Mayara Monteiro Moreira,^a Luanna Monteiro Rodrigues,^a Lidilhone Hamerski^a and Luzineide Wanderley Tinoco*^a

^a Universidade Federal do Rio de Janeiro, Instituto de Pesquisas de Produtos Naturais, Centro de Ciências da Saúde, 21941-902 Rio de Janeiro, RJ, Brasil

^b Universidade Federal do Rio de Janeiro, Laboratório de Avaliação e Síntese de Substâncias Bioativas, Centro de Ciências da Saúde, 21941-902 Rio de Janeiro, RJ, Brasil

Table of contents

Figure S1. Result of biological screening of 214 plants crude extracts.	2
Figure S2. NMR spectra of kaempferol 3-O-β-D-xylopyranosyl-(1→2)-α-L-rhamnopyranoside (2).	3
Figure S3. HSQC NMR spectrum of tricetin-4'-O-methyl flavone (3).	4
Figure S4. IC ₅₀ curves and values for compounds 1 (A), 2 (B) and 3 (C).	5

Supporting Information




Figure S1. Result of biological screening of 214 plants crude extracts. (Each point on this graph corresponds to the result of a bioassay of one crude extract).

Supporting Information




Figure S2. NMR spectra of kaempferol 3-O- β -D-xylopyranosyl-(1 \rightarrow 2)- α -L-rhamnopyranoside (**2**). A) ^1H ; B) HSQC.

Supporting Information




Figure S3. HSQC NMR spectrum of tricetin-4'-O-methyl flavone (3).

Supporting Information




Figure S4. IC_{50} curves and values for compounds **1** (A), **2** (B) and **3** (C).