

Theoretical and Experimental Studies of Oseltamivir-triazole Based Thermoresponsive Organogel

Sumit Kumar^{1,2,†}, Lidong Wu^{3,4,†}, Neha Sharma^{1,†}, Ayushee¹, Kumar Kaushik⁵, Maria Grishina⁶, Bhupendra S Chhikara⁷, Poonam^{2,*}, Vladimir Potemkin^{6,*}, Brijesh Rathi^{1,*}

¹Laboratory for Translational Chemistry and Drug Discovery, Department of Chemistry, Hansraj College University Enclave, University of Delhi, Delhi 110007 India

²Department of Chemistry, Miranda House, University of Delhi, Delhi 110007 India

³Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

⁴Key Laboratory of Control of Quality and Safety for Aquatic Products, Ministry of Agriculture, Chinese Academy of Fishery Sciences, Beijing 100141, China

⁵Fire Chemistry Group, Centre for Fire, Explosives & Environment Safety, Delhi - 110054

⁶South Ural State University, Laboratory of Computational Modeling of Drugs, 454080, Russia

⁷Department of Chemistry, Aditi Mahavidyalaya, University of Delhi, Bawana, Delhi-110039 India

†These authors have contributed equally

Corresponding Author

Poonam (poonam.chemistry@mirandahouse.ac.in); Vladimir Potemkin
(potemkinva@susu.ru); Brijesh Rathi (brijeshrathi@hrc.du.ac.in)




Fig. S1. ^1H NMR (300 MHz)




Fig. S2. ^{13}C NMR for Compound 1.




Fig. S3. UV-Visible Spectra for compound 1.




Fig. S4. Hexane frequency dependence studies of gelator 1




Fig. S5. Decane frequency dependence studies of gelator 1




Fig. S6. Hexane strain and storage modulus for gelator 1.




Fig. S7. Decane strain and storage modulus for gelator 1.




Fig. S8. Powder XRD of gelator 1.