

Supporting information

Combining antioxidant hydrogels with self-assembled microparticles for multifunctional wound dressings

Mariagrazia di Luca,^{†,a,b,c} Manuela Curcio,^{†,d} Emanuele Valli,^{†,e,f} Giuseppe Cirillo,^{*,d} Florida Voli,^e Maria Eugenia Butini,^{b,c} Annafranca Farfalla,^d Elvira Pantuso,^d Antonella Leggio,^d Fiore Pasquale Nicoletta,^d Arianna Tavanti,^a Francesca Iemma^d and Orazio Vittorio^{e,f,g}

^a. Department of Biology, University of Pisa.

^b. Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Center for Musculoskeletal Surgery, Charitéplatz 1, 10117 Berlin, Germany.

^c. Berlin-Brandenburg Center for Regenerative Therapies, Charité – Universitätsmedizin Berlin, Augustenburger Platz 1, 13353 Berlin, Germany.

^d. Department of Pharmacy, Health and Nutritional Sciences, University of Calabria – 87036 – Rende (CS) – Italy.

^e. Children’s Cancer Institute, Lowy Cancer Research Centre, UNSW Sydney, NSW, Australia 2031.

^f. School of Women’s and Children’s Health, Faculty of Medicine, UNSW Sydney, NSW, Australia 2052.

^g. ARC Centre of Excellence for Convergent BioNano Science and Technology, Australian Centre for NanoMedicine, UNSW Sydney, NSW, Australia 2052.

[†] Equal contribution.

* Corresponding author: Department of Pharmacy, Health and Nutritional Sciences, University of Calabria – 87036 – Rende (CS) – Italy. Tel/Fax 0039 0964493011; E-mail: giuseppe.cirillo@unical.it




Fig. S1. UV-Vis absorption spectra of free and conjugated Cur.




Fig. S2. Swelling profiles of HGC (solid line) and HG (dash line) over time.




Fig. S3. MRC-5 viability after 72 h incubation with pulverized H_G (■) and H_{GC} (●) samples suspended in DMEM + FCS 10%.