

## Supporting Information

# **15H-diindolo[2,3-b:1',2'-lm]carbazole: A Novel Rigid Donor for Highly Efficient Thermally Activated Delayed Fluorescence Emitters**


Rajendra Kumar Konidena<sup>a</sup>, Kyung Hyung Lee<sup>a</sup> and Jun Yeob Lee<sup>a\*</sup>, Wan Pyo Hong<sup>b\*</sup>,

<sup>a</sup>School of Chemical Engineering, Sungkyunkwan University, 2066 Seobu-ro, Jangan-gu,  
Suwon, Gyeonggi 440-746, Korea


<sup>b</sup>LG Chem, Ltd, LG Science Park, 30, Magokjungang 10-ro, Gangseo-gu, Seoul, 07796,  
Republic of Korea

Email: [leej17@skku.edu](mailto:leej17@skku.edu), [whongw@lgchem.com](mailto:whongw@lgchem.com)


<b>Fig. S1</b>	Emission spectra of the compounds a) <b>DICzCNTrz</b> and b) <b>CzCNTrz</b> recorded in different solvents.	S2
<b>Fig. S2</b>	Transient PL fluorescence decay (prompt) of the doped films	S2
<b>Fig. S3</b>	TGA traces observed for the compounds	S3
<b>Fig. S4</b>	Electrochemical properties of a) <b>DICzCNTrz</b> and b) <b>CzCNTrz</b> .	S3
<b>Fig. S5</b>	Cyclicvoltammogram of <b>DICz</b>	S4
<b>Fig. S6</b>	a) absorption and b) emission and phosphorescence spectra of <b>DICz</b> donor in THF solution	S4
<b>Fig. S7</b>	<sup>1</sup> H NMR spectra of <b>2</b> recorded in CDCl <sub>3</sub>	S5
<b>Fig. S8</b>	<sup>1</sup> H NMR spectra of <b>3</b> recorded in CDCl <sub>3</sub>	S5
<b>Fig. S9</b>	<sup>1</sup> H NMR spectra of <b>DICz</b> recorded in CDCl <sub>3</sub>	S6
<b>Fig. S10</b>	<sup>1</sup> H NMR spectra of <b>DICzCNTrz</b> recorded in CDCl <sub>3</sub>	S6
<b>Fig. S11</b>	<sup>13</sup> C NMR spectra of <b>DICzCNTrz</b> recorded in CDCl <sub>3</sub>	S7
<b>Fig. S12</b>	<sup>1</sup> H NMR spectra of <b>CzCNTrz</b> recorded in CDCl <sub>3</sub>	S7
<b>Fig. S13</b>	<sup>13</sup> C NMR spectra of <b>CzCNTrz</b> recorded in CDCl <sub>3</sub>	S8




**Fig. S1** Emission spectra of the compounds a) **DICzCNTrz** and b) **CzCNTrz** recorded in different solvents.




**Fig. S2** Transient PL fluorescence decay (prompt) of the doped films in DPEPO.




**Fig. S3** TGA traces observed for the compounds





**Fig. S4** Electrochemical properties of a) DICzCNTrz and b) CzCNTrz.





**Fig. S5** Cyclicvoltammogram of **DICz**




**Fig. S6** a) absorption and b) emission and phosphorescence spectra of DICz donor in THF solution.







**Fig. S10**  $^1\text{H}$  NMR spectra of **DICzCNTrz** recorded in  $\text{CDCl}_3$



**Fig. S11**  $^{13}\text{C}$  NMR spectra of DICzCNTrz recorded in  $\text{CDCl}_3$



**Fig. S12**  $^1\text{H}$  NMR spectra of **CzCNTrz** recorded in  $\text{CDCl}_3$



**Fig. S13**  $^{13}\text{C}$  NMR spectra of **CzCNTrz** recorded in  $\text{CDCl}_3$