

Supporting Information:

Reply to comment on “Zero-thermal-quenching and photoluminescence tuning with the assistance of carriers from defect cluster traps”

Yali Chen,^a Binxun Yu,^{*a} Jing Gou ^{*a} and Shengzhong Frank Liu ^{ab}

^aKey Laboratory of Applied Surface and Colloid Chemistry, Ministry of Education; Shaanxi Key Laboratory for Advanced Energy Devices; Shaanxi Engineering Lab for Advanced Energy Technology; School of Materials Science and Engineering, School of Chemistry and Chemical Engineering, Shaanxi Normal University, Xi'an 710119, China. E-mail: yubx@snnu.edu.cn, goujing@snnu.edu.cn

^bDalian National Laboratory for Clean Energy; iChEM, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, 116023, China; University of the Chinese Academy of Sciences, Beijing 100039, China.




Fig. S1 The XRD patterns of SZ_{1-x}SPO:12%Tb³⁺ (0 ≤ x ≤ 0.6%).

The XRD patterns of SZ_{1-x}SPO:12%Tb³⁺ (0 ≤ x ≤ 0.6%) were presented in Fig. S1 in comparison with the Powder Diffraction File (PDF) card No. 54-1186. All the phosphors were certified as pure phase.