

## Superconductivity in predicted two dimensional $\text{XB}_6$ ( $\text{X} = \text{Ga}, \text{In}$ )

Luo Yan<sup>abcd</sup>, Tao Bo<sup>e</sup>, Peng-Fei Liu<sup>ab</sup>, Liujiang Zhou<sup>f</sup>, Junrong Zhang<sup>ab</sup>, Ming-Hua

Tang<sup>cd</sup>, Yong-Guang Xiao<sup>\*cd</sup> and Bao-Tian Wang<sup>\*abg</sup>

<sup>a</sup> Institute of High Energy Physics, Chinese Academy of Sciences (CAS), Beijing 100049, China. E-mail: wangbt@ihep.ac.cn

<sup>b</sup> Spallation Neutron Source Science Center, Dongguan 523808, China.

<sup>c</sup> Key Laboratory of Key Film Materials & Application for Equipments (Hunan Province), School of Material Sciences and Engineering, Xiangtan University, Xiangtan, Hunan, 411105, China. E-mail: ygxiao@xtu.edu.cn

<sup>d</sup> Hunan Provincial Key Laboratory of Thin Film Materials and Devices, School of Material Sciences and Engineering, Xiangtan University, China.

<sup>e</sup> Songshan Lake Materials Laboratory, Dongguan, Guangdong, 523808, China.

<sup>f</sup> Institute of Fundamental and Frontier Sciences, University of Electronic Science and Technology of China, Chengdu 610054, China.

<sup>g</sup> Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, Shanxi 030006, China.




Figure S1 (a)Top and (b)side views for the rectangular  $\text{XB}_6$  ( $\text{X} = \text{Ga}, \text{In}$ ) with the second minimum energy. The B, Ga(In) atoms are denoted by red and silvery spheres, respectively.