

Hydrogen Bonding Effects on Photocurrent Generation in Porphyrin-Fullerene Photoelectrochemical Devices

Hiroshi Imahori,^{*ab} Jia-Cheng Liu,^a Kohei Hosomizu,^a Tomoo Sato,^c Yukie Mori,^b
Hiroki Hotta,^b Yoshihiro Matano,^{*a} Yasuyuki Araki,^d Osamu Ito,^{*d} Norihiko
Maruyama,^e and Shizuo Fujita^e

^a Department of Molecular Engineering, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan. Fax: 81-75-383-2571; E-mail: imahori@scl.kyoto-u.ac.jp


^b PRESTO, Japan Science and Technology Agency (JST), Japan

^c Department of Chemistry, Graduate School of Pure and Applied Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan


^d Institute of Multidisciplinary Research for Advanced Materials, Tohoku University, CREST, JST, Katahira, Aoba-ku, Sendai 980-8577, Japan

^e International Innovation Center, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan

Expanded Spectra of Figure 2 in the regions of Soret and Q bands (S1):



IR spectra (S2):



Measured in KBr pellet.

AFM images Supplementary Information (S3):

