

## Organocatalytic asymmetric cyanation of isatin derived *N*-Boc ketoimines

Yun-Lin Liu and Jian Zhou\*

Shanghai Key Laboratory of Green Chemistry and Chemical Processes, Department of Chemistry,  
East China Normal University, 3663N Zhongshan Road, Shanghai 200062, China,


E-mail: [jzhou@chem.ecnu.edu.cn](mailto:jzhou@chem.ecnu.edu.cn)


### Supporting Information


#### (Part II)


Content	Page
<b><sup>1</sup>H and <sup>13</sup>C NMR spectra of product 9a-f</b>	<b>2-37</b>
<b><sup>1</sup>H and <sup>13</sup>C NMR spectra of product 10, 13, 14b, 8c, 9c, spirohydantoin I</b>	<b>38-51</b>


<sup>1</sup>H, <sup>13</sup>C NMR and <sup>19</sup>F NMR spectra were recorded on 400 MHz. Chemical shifts are reported in ppm from CDCl<sub>3</sub>, (CD<sub>3</sub>)<sub>2</sub>SO or D<sub>2</sub>O with the solvent resonance as the internal standard. The following abbreviations were used to designate chemical shift multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, h = heptet, m = multiplet, br = broad. Coupling constants, *J*, are reported in Hertz.




































































lyl-11-146-1 c

