

<b>Composition</b>	<b>Mass% Urea</b>	<b>Mass% Glutaric Acid</b>	<b>Mass% Water</b>
1	53.9	1.2	44.9
2	60.58	4.12	35.3
3	10.22	31.04	58.74
4	11.82	38.99	49.19
5	3.72	54.8	41.48
6	52.62	3.58	43.8
7	35.65	5.01	59.34
8	21.6	6.48	71.92
9	20.64	10.65	68.71
10	16.96	12.44	70.6
11	15.97	17.56	66.47
12	15.09	22.12	62.79
13	11.98	21.56	66.46
14	11.43	25.14	63.43
15	11.07	27.47	61.46
16	5.6	52.54	41.86
17	6.18	48.21	45.61
18	7.22	47.67	45.11
19	9.1	46.71	44.19
20	9.71	43.38	46.91
21	10.94	42.79	46.27
22	0.61	66.06	33.33
23	1.24	65.64	33.12

Ternary compositions used to identify all the different solid phase regions in the urea-glutaric acid-water ternary isotherm at 25°C.

<b>Mass% Urea</b>	<b>Mass% Glutaric acid</b>	<b>Mass% Water</b>
8.1	30.7	61.2
7.9	39.3	52.8
7.5	31.7	60.8
7.5	35.9	56.7
7.4	33.7	58.9
5.7	44.5	49.8
5.2	46	48.8
4.9	48	47.1
49.8	0.5	49.7
34	3.2	62.8
18.7	3.2	78.1
14.9	4.3	80.8
11	5.9	83.1
8.8	9.7	81.5
8.3	14.7	77
7.8	17	75.2
7.7	21	71.3
0.8	54.1	45.1
1.6	54.2	44.2

Liquidus compositions for the urea-glutaric acid-water ternary isotherm at 25°C.