

ELECTRONIC SUPPLEMENTARY INFORMATION

Molecular recognition of bisphenol A and its derivatives using *p*-benzoquinone molecule

Noriaki Iwase,^a Takafumi Kinuta,^a Nobuo Tajima,^b Tomohiro Sato,^c Reiko Kuroda,^{c,d} Yoshio Matsubara,^{a*} and Yoshitane Imai^{a*}

^a *Department of Applied Chemistry, Faculty of Science and Engineering, Kinki University, 3-4-1 Kowakae, Higashi-Osaka, Osaka 577-8502, Japan.*

^b *Graduate School of Pure and Applied Sciences, Tsukuba University, 1-1-1 Tenodai, Tsukuba, Ibaraki 305-8571, Japan.*

^c *JST ERATO-SORST Kuroda Chiromorphology Team, 4-7-6, Komaba, Meguro-ku, Tokyo 153-0041, Japan.*

^d *Department of Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8902, Japan.*

Correspondence to: y-imai@apch.kindai.ac.jp, y-matsu@apch.kindai.ac.jp




Fig. ESI-1 Optical absorption spectra (the Kubelka-Munk functions derived from the DRS) of CT complexes **I** (black), **II** (green), and **III** (red).