Supplementary Information

Anion-Controlled Conformational Variation: Structural Modulation and Anion Exchange of Ag(I) Coordination Networks

Wei Wei,^{*a,b*} Hongtao Yu,^{*a*} Feilong Jiang,^{*b*} Bing Liu,^{*a*} Jie Ma,^{*a*} Maochun Hong^{*,*b*}

[a] Department of Chemistry, Capital Normal University, Beijing, 100037, P. R. China.

[b] State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of

Matter, Chinese Academy of Sciences, Fujian, Fuzhou, 350002, P. R. China.

Tel: 86-591-83792460; Fax: +86 591 / 8379 4946

E-mail: hmc@fjirsm.ac.cn.




Fig. S1. The asymmetric unit and the silver (I) coordination environment in **1a**. Symmetry Code: A x, 1-y, z; B -x, 1-y, -z; C -x, y, -z; D -x, -y, -z-1; E -x, 1-y, -z-1; F x, y, z-1; G x, 1-y, z-1.




Fig. S2. The asymmetric unit and the silver (I) coordination environment in **2**. Symmetry Code: A 1-x, y, z; B 2-x, y, z; C 1-x, 1-y, 1/2+z; D x, 1-y, 1/2+z.




Fig. S3. X-ray powder diffraction patterns of 2: a) 2 simulated, b) 2 measured.




Fig. S4. IR (KBr pellet) change procedure during the counteranion exchange of 1a (1) with NH₄ClO₄ after 24 h (2) at 60 °C.




Fig. S5. IR (KBr pellet) change procedure during the counteranion exchange of 1a (1) with NH₄PF₆ after 24 h (2) at 60 °C.





Fig. S6. X-ray powder diffraction patterns of the solid after anion-exchange: a) **1** simulated, b) **1** with NH_4PF_6 , c) **1a** with NH_4PF_6 , d) **1a** with $NaClO_4$ and e) **1a** simulated.






Fig. S7. AFM images of the surface of a crystal of **1**: before anion exchange (top); 15 min after addition of NaNO₂ solution (0.1 M) (middle); 5 h after addition of NaNO₂ solution (0.1 M) (bottom).