1: 1 and 2: 1 Co-Crystals of Alkoxystilbazoles with Tetrafluoroiodobenzenes: Halogen Bonding, a Rare C_{arene}-H···N Hydrogen Bond and Unsymmetric Iodine···Pyridine Interactions

Carsten Präsang, Linda J. McAllister, Adrian C. Whitwood and Duncan W. Bruce*

Department of Chemistry University of York Heslington YORK YO10 5DD (UK) Tel: (+44) 1904 324085

E-mail: duncan.bruce@york.ac.uk



Figure S1 Alternating motif in a side-on view of the 2,3,4,5-tetrafluoroiodobenzene rings in 1.




Figure S2 Upper (coloured green) and lower layer (normal elemental colours) of 2,3,4,5-tetrafluoroiodobenzene chains in successive layers of the structure in 1.




Figure S3 Overlay of the stilbazole with two molecules of 2,3,4,5-tetrafluoroiodobenzene in 1.




Figure S4 Space-filling model of complex 2 showing the back-to-back arrangement of the fluoroarenes.




Figure S5 Halogen-bonded complex 3 showing the equal-probability twisted conformer of the stilbazole.




Figure S6 Complex 3 showing the absence of intermolecular overlap in the stacking direction.




Figure S7 Space-filling model of the packing in complex **3** showing how approach of the alkoxy chain of the stilbazole to the fluoroarene is moderated by a steric interaction.




Fig. S8 Side view of of complex 4.




Fig. S9 Looking down on the superposition of two layers of complex 6.




Figure S10 Diagrams to show the arrangement in complex 7 (recall that in reality the tetrafluoroarene ring is disordered so that it is not clear which is the iodine and which the hydrogen end.