Electronic Supplementary Information: Probing the neutral grapheneionic liquid interface: insights from molecular dynamics simulations[†]

Maxim V Fedorov^{*a} and R. M. Lynden-Bell^{*b‡}

Received Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX First published on the web Xth XXXXXXXX 200X DOI: 10.1039/b000000x

The Electronic Supplementary Information contains: (i) graphical representation of the geometrical setup of the simulation box; (ii) plot of the electrostatic potential in z-direction near the wall.

1 Geometry of the Simulation Box



Fig. 1 Geometrical setup of the simulation box.

^aMax Planck Institute for Mathematics in the Sciences, Inselstrasse 22, Leipzig 04103, Germany. Fax: +49 341 9959 999; Tel: +49 341 9959 804; E-mail: fedorov@mis.mpg.de

^bUniversity of Cambridge, University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW. Tel: +44 1223 336535 E-mail: rmlb@cam.ac.uk

2 Electrostatic Potential at the Wall



Fig. 2 The interfacial number densities of [dmim][Cl] cations (red) and anions (blue). The electrostatic potential in the direction perpeidicular to the wall (z-direction) is shown by black below the density curves (the corresponding ordinate is on the right side of the plot).