

Electronic Supplementary Material: First example of a high-level correlated calculation of the indirect spin-spin coupling constants involving tellurium: tellurophene and divinyl telluride

Yury Yu. Rusakov,^a Leonid B. Krivdin,^{*a} Freja F. Østerstrøm,^b Stephan P. A. Sauer,^b Vladimir A.
5 Potapov,^a Svetlana V. Amosova^a

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Table 1 Exponents, taken from "relativistic" Dyll's basis set dyall.av3z,^{1,2} and contraction coefficients of the av3z-J basis set for tellurium

[illegible]

	5.9722776		1.0																	
	3.1602785			1.0																
	1.6294969						1.0													
	0.81785307						1.0													
	0.38849309							1.0												
	0.15854467								1.0											
	0.06274744								1.0											
<i>f</i>	0.14440990	1.0																		
	0.34378542		1.0																	

^a Footnote text.

Notes and references

^a A.E. Favorsky Irkutsk Institute of Chemistry, Russian Academy of Sciences, Siberian Branch, 1 Favorsky St., 664033 Irkutsk, Russia. E-mail: krivdin_office@irioch.irk.ru

^b Department of Chemistry, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark.

1 K. G. Dyall, *Theor. Chem. Acc.* 2002, **108**, 335.

2 K. G. Dyall, *Theor. Chem. Acc.* 2006, **115**, 441.