a	. •		•	•
Suppo	rtıng	ınt	ormat	10n

 ${\bf Liquid\hbox{-}liquid\hbox{-}extraction\hbox{ of neodymium}(III)\hbox{ by dialkylphosphate ionic liquids from a cidic medium: the importance of the ionic liquid cation}$ 

Alok Rout, Justyna Kotlarska, Wim Dehaen, Koen Binnemans\*




Figure S1.  $^{1}H$  NMR spectrum of [C<sub>6</sub>mim][DEHP].




Figure S2. <sup>13</sup>C NMR spectrum of [C<sub>6</sub>mim][DEHP].




Figure S3. <sup>31</sup>P NMR spectrum of [C<sub>6</sub>mim][DEHP].




Figure S4. <sup>1</sup>H NMR spectrum of [N<sub>4444</sub>][DEHP].




Figure S5.  $^{13}$ C NMR spectrum of [N<sub>4444</sub>][DEHP].




Figure S6. <sup>31</sup>P NMR spectrum of [N<sub>4444</sub>][DEHP].




Figure S7. <sup>1</sup>H NMR spectrum of [C<sub>6</sub>mpyr][DEHP].




Figure S8. <sup>13</sup>C NMR spectrum of [C<sub>6</sub>mpyr][DEHP].




Figure S9. <sup>31</sup>P NMR spectrum of [C<sub>6</sub>mpyr][DEHP].