

Supplementary Material

DPT tautomerisation of the $G \cdot A_{\text{syn}}$ and $A^* \cdot G^*_{\text{syn}}$ DNA mismatches: A QM/QTAIM combined atomistic investigation

Ol'ha O. Brovarets' and Dmytro M. Hovorun[✉]

^aDepartment of Molecular and Quantum Biophysics, Institute of Molecular Biology and Genetics, National Academy of Sciences of Ukraine, 150 Akademika Zabolotnoho Str., 03680 Kyiv, Ukraine

^bDepartment of Molecular Biotechnology and Bioinformatics, Institute of High Technologies, Taras Shevchenko National University of Kyiv, 2-h Akademika Hlushkova Ave., 03022 Kyiv, Ukraine

[✉]Corresponding author. Email: dhovorun@imbg.org.ua.




Fig. S1 Profiles of: (a) the electronic energy E of the base mispair and (b) the first derivative of the electronic energy E with respect to the IRC - $dE/dIRC$ - along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*. The position of the $TS_{A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}}$ corresponds to IRC=0.00 Bohr.




Fig. S2 Profile of the dipole moment μ of the base mispair along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*.




Fig. S3 Profiles of: **(a)** the electron density ρ ; **(b)** the Laplacian of the electron density $\Delta\rho$; **(c)** the ellipticity ϵ and **(d)** the energy of the H-bond E_{HB} , estimated by the EML formula [1,2], at the (3,-1) BCPs of the covalent and hydrogen bonds along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*.




Fig. S4 Profiles of: **(a)** the distance $d_{A...B}$ between the electronegative A and B atoms; **(b)** the distance $d_{AH/HB}$ between the hydrogen and electronegative A or B atoms and **(c)** the angle $\angle AH...B$ of the $AH...B$ H-bonds along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*.




Fig. S5 Profiles of the NBO charges of the hydrogen atoms involved in the $O6H_I \cdots N6/N6H_I \cdots O6$ and $N1H_{II} \cdots N7/N7H_{II} \cdots N1$ H-bonds along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*.




Fig. S6 Profiles of: (a) the distance $R(H-H)$ between the glycosidic hydrogens and (b) the α_1 ($\angle N9H(A)H(G_{syn})$) and α_2 ($\angle N9H(G_{syn})H(A)$) glycosidic angles along the IRC of the $A^* \cdot G^*_{syn} \leftrightarrow A \cdot G^*_{syn}$ tautomerisation *via* the DPT obtained at the B3LYP/6-311++G(d,p) level of theory *in vacuo*.

References.

1. Espinosa E., Molins E., Lecomte C. Hydrogen bond strengths revealed by topological analyses of experimentally observed electron densities, *Chem. Phys. Lett.*, 1998, **285**, 170–173.
2. Mata I., Alkorta I., Espinosa E., Molins E. Relationships between interaction energy, intermolecular distance and electron density properties in hydrogen bonded complexes under external electric fields, *Chem. Phys. Lett.*, 2011, **507**, 185–189.