

Lyotropic Liquid Crystal Phases of Phytantriol in a Protic Ionic Liquid with Fluorous Anion

Yan Shen,^{a,b} Tamar L. Greaves,^{a,d} Danielle F. Kennedy,^a Asoka Weerawardena,^{a,d} Nigel Kirby,^c Gonghua Song,^b and Calum J. Drummond^{a,d*}


^a CSIRO Materials Science and Engineering (CMSE), Bag 10, Clayton, Vic. 3169, Australia.

^b Shanghai Key Laboratory of Chemical Biology, School of Pharmacy, East China University of Science and Technology, Shanghai 200237, PR China.

^c Australian Synchrotron, 800 Blackburn Road, Clayton, Vic. 3168, Australia.

^d School of Applied Sciences, College of Science, Engineering and Health, RMIT University, GPO Box 2476, Melbourne, Victoria 3001, Australia.

*Corresponding author. Email: calum.drummond@rmit.edu.au











Figure S1. Stackplots of SAXS/WAXS patterns at 25 °C for phytantriol in a) MImOF (no water), b) MImOF90wt%-water10wt%, c) MImOF75wt%-water25wt%, d) MImOF50wt%-water50wt%, and e) MImOF25wt%-water75wt%. Concentrations refer to the proportion of phytantriol in the binary or tertiary system.








Figure S2. Stackplots of IR spectra at 20 °C for phytantriol in a) MImOF (no water), b) MImOF90wt%-water10wt%, c) MImOF75wt%-water25wt%, d) MImOF50wt%-water50wt% and e) MImOF25wt%-water75wt%.




Figure S3. IR spectra at 20 °C for phytantriol concentrations between 30 and 80 wt% in the solvent mixture of 75 wt% MImOF and 25 wt% water.




Figure S4. IR spectra at 20 °C for phytantriol concentrations between 30 and 80 wt% in the solvent mixture of 50 wt% MImOF and 50 wt% water.