

Electronic Supplementary Material

Symmetrical and Unsymmetrical Bronsted acidic ionic liquids for the effective conversion of Fructose to 5-Hydroxymethyl furfural

Deepali A. Kotadia, Saurabh S. Soni*

1. Characterization of Ionic liquids


- ^1H and ^{13}C NMR of [MMIM] HSO_4 (Figure S1)
- ^1H and ^{13}C NMR of [MMBIM] HSO_4 (Figure S2)
- ^1H and ^{13}C NMR of [HMBIM] HSO_4 (Figure S3)
- ^1H and ^{13}C NMR of [PSMBIM] HSO_4 (Figure S4)
- TGA of [MMIM] HSO_4 (Figure S5)
- TGA of [MMBIM] HSO_4 (Figure S6)
- TGA of [PSMBIM] HSO_4 (Figure S7)
- TGA of [HMBIM] HSO_4 (Figure S8)

2. Identification and Characterization of 5-Hydroxymethylfurfural


- GC-MS of 5-Hydroxymethyl furfural (Figure S9)
- HPLC measurement using UV and RI detector (Figure S10)
- ^1H and ^{13}C NMR of HMF (Figure S11)

The ^1H and ^{13}C NMR of synthesized Ionic liquids


^1H NMR of [MMIM] HSO_4^- (Figure S1)




^{13}C NMR of [MMIM] HSO_4^-




¹H NMR of [MMBIM] HSO₄ (Figure S2)




¹³C NMR of [MMBIM] HSO₄




¹H NMR of [HMBIM] HSO₄ (Figure S3)




¹³C NMR of [HMBIM] HSO₄



¹H NMR of [PSMBIM] HSO₄ (Figure S4)




¹³C NMR of [PSMBIM] HSO₄




Thermogravimetric analysis graph


[MMIM] HSO_4 (Figure S5)




[MMBIM] HSO_4 (Figure S6)



[HMBIM] HSO_4 (Figure S7)



[PSMBIM] HSO_4 (Figure S8)




Identification and Characterization of 5-Hydroxymethylfurfural


GC-MS of 5-Hydroxymethyl furfural (**Figure S9**)




HPLC-UV of 5-Hydroxymethyl furfural (**Figure S10**)



HPLC-RI of 5-Hydroxymethyl furfural



¹H NMR of 5-HMF (Figure S11)



¹³C NMR of 5-HMF

