

Electronic Supplementary Information

Mesoporous Zr-SBA-16 catalysts for glycerol valorization processes: towards biorenewable formulations

Camino Gonzalez-Arellano^{a*}, Leticia Parra-Rodriguez^a, Rafael Luque^b

^a*Departamento de Química Orgánica y Química Inorgánica, Universidad de Alcalá, Facultad de Farmacia, Autovía A2, 33.600, 28871, Alcalá de Henares, Madrid (Spain) E-mail: camino.gonzalez@uah.es, Tel.: +34-918854677; Fax: +34-918854683.*

^b*Departamento de Química Orgánica, Universidad de Córdoba, Campus de Rabanales, Edificio Marie Curie (C-3), Ctra Nnal IV-A, Km 396, 14014 Córdoba (Spain), E-mail: q62alsor@uco.es*




Figure 1S. Chromatogram reaction with Zr-SBA-16(100), 1 mmol glycerol, 5 mmol levulinic acid, 15h, 100°C (selectivity 71/29/0).




Figure 2S. Chromatogram reaction with Zr-SBA-16(100), 1 mmol glycerol, 5 mmol levulinic acid, 8h, 140°C (selectivity 11/64/25).




Figure 3S. Chromatogram reaction with Zr-SBA-16(100), 1 mmol glycerol, 5 mmol levulinic acid, 15h, 140°C (selectivity 0/50/50).