Supporting Information

Fast and precise quantification of serum biomarkers and simultaneous recognition of multiple diseases enabled by stable isotope-labelled peptides assistant high-throughput MRM strategy

Anqi Hu ^{1‡}, Zhenxin Wang^{1,2‡}, Juanjuan Xie¹, Ling Lin¹, Pengyuan Yang^{1,3,4,5} Lei Zhang^{1*}, Huali Shen^{1,4,5*}, and Yi-ou Cao^{1*}

¹ Minhang Hospital and Institutes of Biomedical Sciences, Fudan University, Shanghai, 201199, China

² Laboratory Medicine of Zhongshan Hospital, Fudan University, Shanghai, 200032, China

³ Department of Chemistry, Fudan University, Shanghai, 200032, China

⁴ Department of Systems Biology for Medicine and School of Basic Medical Sciences, Fudan University, Shanghai, China

⁵ NHC Key Laboratory of Glycoconjugates Research, Fudan University, Shanghai 200032, China

[‡] These authors contributed equally to this work.

* Corresponding authors

Lei Zhang (zhanglei123@fudan.edu.cn)

Huali Shen (shenhuali@fudan.edu.cn)

Yi-ou Cao (yioucao_doctor@126.com)

Contents

- 1, supporting figures
- 2、 supporting tables

Supporting Figures



Fig. S1 The molecular types of the 500-serum proteins corresponding to the SIL peptides panel.



Fig. S2 Estimated concentration distribution of 113 proteins in all samples(A). The distribution of average L/H ratio in each group (B) and average L/H ratio of representative protein in each group(C).



Fig.S3 Venn diagram displaying the DEPs summary statistics(A). Top 10 terms of enriched biological processes of DEPs in AS (B) and cancer (C). Cancer refers to LC and BC in our study. DEPs, differentially expressed proteins.