## **Supplementary Information**

Barcode signal amplifying strategy for sensitive and accurate protein detection on LC-MS/MS

Chang Shu<sup>1,2</sup>, Teng-Fei Li<sup>3</sup>, Duo Li<sup>2</sup>, Zhong-Qiu Li<sup>1</sup>, Xing-Hua Xia<sup>1\*</sup>

\*Corresponding author E-mail address: xhxia@nju.edu.cn.

- <sup>1</sup> State Key Laboratory of Analytical Chemistry for Life Science, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing, 210023, China.
- <sup>2</sup> Department of Pharmaceutical Analysis, School of Pharmacy, China Pharmaceutical University, Nanjing, 210009, China.
- <sup>3</sup>School of Pharmacy, Department of Clinical Pharmacology, Sir Run Run Hospital, Nanjing Medical University, Nanjing 211166, China



Figure S-1. Synthetic route of compounds



Figure S-2. The emission spectra of QDs, Fmoc-ss@QDs and Apt15@ss@QDs solution.



Figure S-3. The FTIR spectra of Apt15 (a), Apt15@QDs (b) and Apt15@ss@QDs (c).



Figure S-4. The FTIR spectra of Apt29 (a) and Apt29@MNPs (b).



**Figure S-5.** The effects of different amounts of Tris-HCl solution (A), Apt29@MB (B), Apt15@ss@QDs (C) and TCEP (D) on thrombin detection.



Figure S-6. LC-MS/MS spectra of mass barcode and internal standard



Figure S-7. High resolution mass spectrum of Fmoc-ss compound



Figure S-8. <sup>1</sup>H NMR of Fmoc-ss compound



Figure S-9. High resolution mass spectrum of Fmoc-R-ss compound



Figure S-10. <sup>1</sup>H-NMR of Fmoc-R-ss compound



Figure S-11. High resolution mass spectrum of mass barcode compound



Figure S-12. <sup>1</sup>H-NMR of mass barcode compound