

Supplementary Information

A crystalline solid adduct of sulfathiazole-amantadine: The first dual-drug molecular salt containing both antiviral and antibacterial ingredients

Ling-Yang Wang^a, Yue-Ming Yu^a, Ming-Chao Yu^a, Yan-Tuan Li^{a,b*}, Zhi-Yong Wu^a, Cui-Wei Yan^{a,*}

^a*School of Medicine and Pharmacy and College of Marine Life Science, Ocean University of China, Qingdao, Shandong 266003, PR China.*

^b*Laboratory for Marine Drugs and Bioproducts, Qingdao National Laboratory for Marine Science and Technology, 266003, PR China.*

* Correspondence to: Yan-Tuan Li and Cui-Wei Yan.

E-mail addresses: yantuanli@ouc.edu.cn (Yan-Tuan Li), cuiweiyan@ouc.edu.cn (Cui-Wei Yan).

This ESI for *CrystEngComm*, 2020, **22**, 3804-3813, DOI: 10.1039/D0CE00368A, first published on 7th May 2020, was updated on 8th June 2023.

*Correspondence to: Yan-Tuan Li and Cui-Wei Yan.

E-mail address: yantuanli@ouc.edu.cn; Cui-Wei Yan, E-mail: cuiweiyan@ouc.edu.cn.

Contents

Fig. S1. FT-IR spectra of SFZ-ATD molecular salt compared with its starting materials.

Fig. S2. XRPD comparison of simulated pattern for SFZ-ATD molecular salt and the residual solids after the solubility measurements.

Fig. S3. IDR profiles of pure SFZ, ATD and SFZ-ATD molecular salt in different media.




Fig. S1. FT-IR spectra of SFZ-ATD molecular salt compared with its starting materials.
(a) ATD, (b) SFZ and (c) SFZ-ATD.




Fig. S2. XRPD comparison of simulated pattern for SFZ-ATD molecular salt and the residual solids after the solubility measurements.




Fig. S3. IDR profiles of pure SFZ, ATD and SFZ-ATD molecular salt in different media.