

Ecotoxicological impact of graphene oxide: toxic effects on the model organism *Artemia franciscana*

Cavion Federica^a, Fusco Laura^b, Sosa Silvio^a, Manfrin Chiara^a, Alonso Beatriz^c, Zurutuza Amaia^c, Della Loggia Roberto^b, Tubaro Aurelia^a, Prato Maurizio^{b,d,e}, Pelin Marco^{a*}

*corresponding author

^a Department of Life Sciences, University of Trieste, 34127 Trieste, Italy

^b Department of Chemical and Pharmaceutical Sciences, University of Trieste, 34127 Trieste, Italy

^c Graphenea, 20009 Donostia-San Sebastián, Spain

^d Carbon Nanobiotechnology Laboratory CIC BiomaGUNE, 20009 Donostia-San Sebastián, Spain

^e Basque Foundation for Science (IKERBASQUE), 48013 Bilbao, Spain

Supplementary materials

Figure S1. Physico-chemical characterization of GO. (A) Raman Spectra of the Graphene Oxide deposited in a glass slide; each color one point. (B) HR-TEM image of GO; the e-beam reduced partially the graphene oxide and the honey-comb structure is recovered. (C) SEM image of GO flakes deposited on a SiO₂/Si substrate.




Figure S2. Uptake of GO by *A. franciscana* adults. Representative images of *A. franciscana* adults exposed to GO for 24 h (A) in absence of food and 48 h (B), in the absence and in the presence of food. The black and green arrows indicate GO and food into the lumen of the digestive tract, respectively. Pictures were acquired under the stereomicroscope at a 1X magnification. Scale bars: 1 mm.

