

## Supplementary Material:

# Attosecond laser control of photoelectron angular distributions in XUV-induced ionization of H<sub>2</sub>

Roger Y. Bello,<sup>1</sup> Fernando Martín,<sup>2,3,4,\*</sup> and Alicia Palacios<sup>2,5</sup>

<sup>1</sup>*Lawrence Berkeley National Laboratory,  
Chemical Sciences, Berkeley, California 94720, USA.*

<sup>2</sup>*Departamento de Química, Módulo 13,  
Universidad Autónoma de Madrid, 28049 Madrid, Spain.*

<sup>3</sup>*Instituto Madrileño de Estudios Avanzados en Nanociencia (IMDEA-Nanociencia),  
Cantoblanco, 28049 Madrid, Spain.*

<sup>4</sup>*Condensed Matter Physics Center (IFIMAC),  
Universidad Autónoma de Madrid, 28049 Madrid, Spain.*

<sup>5</sup>*Institute of Advanced Research in Chemical Sciences (IAdChem),  
Universidad Autónoma de Madrid, 28049 Madrid, Spain*

---

\* fernando.martin@uam.es

## MOVIES SHOWING THE VARIATION OF THE ASYMMETRY PARAMETERS WITH PUMP-PROBE DELAY

1. The movie *Video\_diss\_project* shows the variation with pump-probe delay of the asymmetry parameter for XUV pump-IR probe dissociative ionization of H<sub>2</sub> as a function of EKE and NKE when electrons are detected and the proton emission direction. is determined.
2. The movie *Video\_diss\_noproject* shows the variation with pump-probe delay of the asymmetry parameter for XUV pump-IR probe dissociative ionization of H<sub>2</sub> as a function of EKE and NKE when electrons are detected irrespective of the proton emission direction.
3. The movie *Video\_bound* shows the variation with pump-probe delay of the asymmetry parameter for XUV pump-IR probe non dissociative ionization of H<sub>2</sub> as a function of EKE and NKE.