

Supplementary material

Table S1. Dietary energy distribution of the normal-fat diet (NFD, #D12450B) and high-fat diet (HFD, #D12451) purchased from Research Diets Co. Ltd. (New Brunswick, NJ, USA).

Nutritional Components	Normal-fat diet	High-fat diet
	(10% dietary fat)	(45% dietary fat)
Protein	20	20
Carbohydrate	70	35
Fat	10	45
Total kcal/g	100	100

Table S2. Composition of the normal-fat diet (NFD, #D12450B) and high-fat diet (HFD, #D12451) used in the animal experiment.

Ingredients	Normal-fat diet		High-fat diet	
	(10% dietary fat)		(45% dietary fat)	
	g	kcal	g	kcal
Casein, 30 mesh	200	800	200	800
L-cystine	3	12	3	12
Corn starch	315	1260	72.8	291
Maltodextrin 10	35	140	100	400
Sucrose	350	1400	172.8	691
Cellulose, BW200	50	0	50	0
Soybean oil	25	225	25	225
Lard	20	180	177.5	1598
Mineral mix S10026	10	0	10	0
Dicalcium phosphate	13	0	13	0
Calcium carbonate	5.5	0	5.5	0
Potassium citrate	16.5	0	16.5	0
Vitamin mix V10001	10	40	10	40
Choline bitartrate	2	0	2	0
FD&C yellow dye	0.05	0	-	-
FD&C red dye	-	-	0.05	0

Fig. S1. Effects of simvastatin and Ganoderma acid A administration on the food intake in mice fed a high-fat diet.




Fig. S2. Effects of simvastatin and Ganoderma acid A administration on the adipocyte size in mice fed a high-fat diet.




Fig. S3. Effects of simvastatin and Ganoderma acid A administration on liver lipid accumulation in mice fed a high-fat diet.

