

Supplementary information

Complementary techniques to analyse pericellular matrix formation by human MSC within hyaluronic acid hydrogels

Christoph Salzlechner¹†, Anders Runge Walther^{1,2}†, Sophie Schell^{1,3}, Nicholas Groth Merrild¹, Tabasom Haghighi¹, Isabella Huebscher¹, Gerhard Undt⁴, Kathleen Fan⁵, Mads Bergholt¹, Martin Hedegaard², and Eileen Gentleman^{1}*

¹ Centre for Craniofacial and Regenerative Biology, King's College London, London SE1 9RT, United Kingdom

² Department of Chemical Engineering, Biotechnology and Environmental Technology, University of Southern Denmark, DK-5230 Odense, Denmark

³ Department of Conservative Dentistry, Centre of Dentistry, Oral Medicine and Maxillofacial Surgery, University Hospital Tübingen, 72076 Tübingen, Germany

⁴ University Clinic of Dentistry, Department of Oral Surgery, Medical University of Vienna, Sensengasse 2a, 1090 Vienna, Austria

⁵ Department of Oral and Maxillofacial Surgery, King's College Hospital, London, SE5 9RS, United Kingdom

†These authors contributed equally

*To whom correspondence should be addressed: eileen.gentleman@kcl.ac.uk

Highest academic title: Dr Christoph Salzlechner, Prof Gerhard Undt, Dr Sophie Schell, Dr Mads Bergholt, Dr Martin Hedegaard, Dr Kathleen Fan, Dr Eileen Gentleman




Figure S1 Representative confocal images of hydrogel encapsulated MSCs. **(A)** Fluorescent non-canonical amino acid tagging (red) and nucleus (blue). **(B)** Differential interference contrast (DIC) images of the cells shown in **A** for visualization of cell outline and **(C)** merged view of red and DIC channels.